



Chapter 4: Phonons I

Crystal Vibrations





OUTLINES

- ☐ *Vibrations of crystals with monatomic basis*
- ☐ *Two atoms per primitive basis*
- ☐ *Quantization of elastic waves*
- ☐ *Phonon momentum*
- ☐ *Elastic scattering of phonons*


Major Elementary Excitation in Solids

	Name	Field
	Electron	—
	Photon	Electromagnetic wave
	Phonon	Elastic wave
	Plasmon	Collective electron wave
	Magnon	Magnetization wave
—	Polaron	Electron + elastic deformation
—	Exciton	Polarization wave

Figure 1 Important elementary excitations in solids.

Displacement of Planes of Atoms in a *Longitudinal* Wave

U_s is defined as the displacement for the plane s from its equilibrium position



Displacement of Planes of Atoms in a *Transverse* Wave




Figure 3 Planes of atoms as displaced during passage of a transverse wave.

Hooke's Law

- ❑ We assume the elastic response of the crystal is a linear function of the forces.
- ❑ The elastic energy is a quadratic function of the relative displacement of any two points in the crystal.
- ❑ **Hooke's Law** : The force exerted on the plane s as caused by the displacement of the plane $s+p$ is directly proportional to the difference of the displacement $u_{s+p} - u_s$. For nearest neighbor interaction, $p = \pm 1$
- ❑ Hence, the total force on plane s from planes $s+1$, and $s-1$ is

$$F_s = C(u_{s+1} - u_s) + C(u_{s-1} - u_s) \quad (1)$$

The equation of motion of the plane s is


$$M \frac{d^2 u_s}{dt^2} = C(u_{s+1} + u_{s-1} - 2u_s) , \quad \text{C: force constant between nearest neighbor planes for one atom in the plane} \quad (2)$$

With time dependence, $u = u \exp(-i\omega t)$

$$-M\omega^2 u_s = C(u_{s+1} + u_{s-1} - 2u_s) \quad (3)$$

By the traveling wave solution for a periodic set of atomic planes with a spacing of "a" , $u_s = u \exp(isKa)$

$$u_{s\pm 1} = u \exp(isKa) \exp(\pm iKa) , \quad (4)$$



$$\begin{aligned}
 & -\omega^2 M u \exp(isKa) \\
 & = Cu \{ \exp[i(s+1)Ka] + \exp[i(s-1)Ka] - 2 \exp(isKa) \} . \quad (5)
 \end{aligned}$$

$$\omega^2 M = -C [\exp(iKa) + \exp(-iKa) - 2] . \quad (6)$$

$$\omega^2 = (2C/M)(1 - \cos Ka) . \quad (7)$$

At the first Brillouin zone boundary, $K = \pi/a$, and $-\pi/a$,

$$d\omega^2/dK = (2Ca/M) \sin Ka = 0 \quad (8)$$

$$\omega^2 = (4C/M) \sin^2 \frac{1}{2} Ka ; \quad \boxed{\omega = (4C/M)^{1/2} |\sin \frac{1}{2} Ka|} . \quad (9)$$

ω vs κ Dispersion for Monoatomic Lattice





Figure 4 Plot of ω versus K . The region of $K \ll 1/a$ or $\lambda \gg a$ corresponds to the continuum approximation; here ω is directly proportional to K .



$$\frac{u_{s+1}}{u_s} = \frac{u \exp[i(s+1)Ka]}{u \exp(isKa)} = \exp(iKa) . \quad \begin{array}{l} -\pi < \mathbf{K} a < \pi \\ -\pi/a < \mathbf{K} < \pi/a \end{array} \quad (10)$$

The meaningful range of $\mathbf{K}$ is only inside the first Brillouin Zone of the linear lattice.

$$u_{s+1}/u_s = \exp(iKa) \equiv \exp(i2\pi n) \exp[i(Ka - 2\pi n)] \equiv \exp(iK'a) , \quad (11)$$

$$\mathbf{K}' = \mathbf{K} - 2n\pi/a = \mathbf{K} - n\mathbf{G}$$

We can always subtract a reciprocal lattice vector $\mathbf{G}$ from $\mathbf{K}$ to become $\mathbf{K}'$, to be inside the first Brillouin zone. “**Reduced zone scheme !**”

At the zone boundary, $K_{max} = \pi/a$, and $-\pi/a$

$$u_s = u \exp(\pm is\pi) = u (-1)^s . \quad (12)$$

This is not a traveling wave, but **a standing wave**; alternating atoms oscillate in opposite phases. $\mathbf{U}_s$ equals to $\mathbf{u}$ or $-\mathbf{u}$, depending on s is an even, or odd integer.

Reciprocal Lattice Vector

To proceed further with the Fourier analysis of the electron concentration we must find the vectors $\mathbf{G}$ of the Fourier sum $\sum n_{\mathbf{G}} \exp(i\mathbf{G} \cdot \mathbf{r})$ as in (9).

We construct the axis vectors $\mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3$ of the reciprocal lattice: 倒晶格

$$\mathbf{b}_1 = 2\pi \frac{\mathbf{a}_2 \times \mathbf{a}_3}{\mathbf{a}_1 \cdot \mathbf{a}_2 \times \mathbf{a}_3} ; \quad \mathbf{b}_2 = 2\pi \frac{\mathbf{a}_3 \times \mathbf{a}_1}{\mathbf{a}_1 \cdot \mathbf{a}_2 \times \mathbf{a}_3} ; \quad \mathbf{b}_3 = 2\pi \frac{\mathbf{a}_1 \times \mathbf{a}_2}{\mathbf{a}_1 \cdot \mathbf{a}_2 \times \mathbf{a}_3} . \quad (13)$$

If $\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3$ are primitive vectors of the crystal lattice, then $\mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3$ are primitive vectors of the reciprocal lattice. Each vector defined by (13) is orthogonal to two axis vectors of the crystal lattice. Thus $\mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3$ have the property

$$\mathbf{b}_i \cdot \mathbf{a}_j = 2\pi \delta_{ij} , \quad (14)$$

where $\delta_{ij} = 1$ if $i = j$ and $\delta_{ij} = 0$ if $i \neq j$.

Points in the reciprocal lattice are mapped by the set of vectors

$$\mathbf{G} = v_1 \mathbf{b}_1 + v_2 \mathbf{b}_2 + v_3 \mathbf{b}_3 , \quad (15)$$

where v_1, v_2, v_3 are integers. A vector $\mathbf{G}$ of this form is a reciprocal lattice vector.

倒晶格向量

Reciprocal Lattice to sc Lattice

Simple Cubic

The primitive translation vectors of a simple cubic lattice may be taken as the set

$$\mathbf{a}_1 = a\hat{\mathbf{x}} ; \quad \mathbf{a}_2 = a\hat{\mathbf{y}} ; \quad \mathbf{a}_3 = a\hat{\mathbf{z}} . \quad (27a)$$

Here $\hat{\mathbf{x}}, \hat{\mathbf{y}}, \hat{\mathbf{z}}$ are orthogonal vectors of unit length. The volume of the cell is $\mathbf{a}_1 \cdot \mathbf{a}_2 \times \mathbf{a}_3 = a^3$. The primitive translation vectors of the reciprocal lattice are found from the standard prescription (13):

$$\mathbf{b}_1 = (2\pi/a)\hat{\mathbf{x}} ; \quad \mathbf{b}_2 = (2\pi/a)\hat{\mathbf{y}} ; \quad \mathbf{b}_3 = (2\pi/a)\hat{\mathbf{z}} . \quad (27b)$$

Here the reciprocal lattice is itself a simple cubic lattice, now of lattice constant $2\pi/a$.

The boundaries of the first Brillouin zones are the planes normal to the six reciprocal lattice vectors $\pm\mathbf{b}_1, \pm\mathbf{b}_2, \pm\mathbf{b}_3$ at their midpoints:

$$\pm\frac{1}{2}\mathbf{b}_1 = \pm(\pi/a)\hat{\mathbf{x}} ; \quad \pm\frac{1}{2}\mathbf{b}_2 = \pm(\pi/a)\hat{\mathbf{y}} ; \quad \pm\frac{1}{2}\mathbf{b}_3 = \pm(\pi/a)\hat{\mathbf{z}} . \quad (28)$$

The six planes bound a cube of edge $2\pi/a$ and of volume $(2\pi/a)^3$; this cube is the first Brillouin zone of the sc crystal lattice.

Group Velocity

The transmission velocity of a wave packet is the group velocity

$$v_g = d\omega/dK ,$$

or

$$\mathbf{v}_g = \text{grad}_{\mathbf{K}} \omega(\mathbf{K}) , \quad (13)$$

From Eq. 9,
$$v_g = (Ca^2/M)^{1/2} \cos \frac{1}{2} Ka . \quad (14)$$

At zone boundary, $K = \pi/a$, $V_g = 0$ for standing wave

At the zone center, $Ka \ll 1$, the continuum approximation

$$\omega^2 = (C/M)K^2a^2 . \quad (15)$$

$$\mathbf{v}_g = (C/M)^{1/2} a \quad V_g \sim \text{is nearly a constant}$$

See Figure 6

Group Velocity v_g vs K of Mono Atomic Lattice




Figure 6 Group velocity v_g versus K , for model of Fig. 4. At the zone boundary $K = \pi/a$ the group velocity is zero.

The Traveling Wave Description of the Atomic Displacement in a linear lattice




Figure 5 The wave represented by the solid curve conveys no information not given by the dashed curve. Only wavelengths longer than $2a$ are needed to represent the motion.

a : lattice spacing

$$\lambda / 2 > a$$

$$\lambda > 2a$$

$$K < \pi/a$$

Derivation of Force Constant from Experiment

For longer range force, we include p nearest planes of contributions to ω

$$\omega^2 = (2/M) \sum_{p>0} C_p (1 - \cos pKa) . \quad (16a)$$

We times $M \cos rKa$ term on both sides, and integrate over K

$$\begin{aligned} M \int_{-\pi/a}^{\pi/a} dK \omega_K^2 \cos rKa &= 2 \sum_{p>0} C_p \int_{-\pi/a}^{\pi/a} dK (1 - \cos pKa) \cos rKa \\ &= -2\pi C_r / a . \end{aligned} \quad (16b)$$

Note the integral vanishes, except for $p = r$, and that equals to $-\pi/a$

$$C_p = -\frac{Ma}{2\pi} \int_{-\pi/a}^{\pi/a} dK \omega_K^2 \cos pKa \quad (17)$$

From experimentally measured ω_K , we will derive C_p

Displacement of a Diatomic Linear Crystal Structure





Figure 9 A diatomic crystal structure with masses M_1 , M_2 connected by force constant C between adjacent planes. The displacements of atoms M_1 are denoted by $u_{s-1}, u_s, u_{s+1}, \dots$, and of atoms M_2 by v_{s-1}, v_s, v_{s+1} . The repeat distance is a in the direction of the wavevector K . The atoms are shown in their undisplaced positions.

Considering only nearest neighbor interaction, force constant C are identical between all pairs of near-neighbor planes.

Equation of Motion for a Diatomic Linear Crystal


$$\begin{aligned} M_1 \frac{d^2 u_s}{dt^2} &= C(v_s + v_{s-1} - 2u_s) ; \\ M_2 \frac{d^2 v_s}{dt^2} &= C(u_{s+1} + u_s - 2v_s) . \end{aligned} \quad (18)$$

Traveling wave solution

$$\begin{cases} u_s = u \exp(isKa) \exp(-i\omega t) ; \\ v_s = v \exp(isKa) \exp(-i\omega t) . \end{cases} \quad (19)$$

a as the distance between nearest identical planes,
but not nearest neighbor planes.

$$\begin{aligned} -\omega^2 M_1 u &= Cv[1 + \exp(-iKa)] - 2Cu ; \\ -\omega^2 M_2 v &= Cu[\exp(iKa) + 1] - 2Cv . \end{aligned} \quad (20)$$

ω vs K for a Diatomic Linear Crystal

Solution exists only if the determinant of the coefficients vanishes

$$\begin{vmatrix} 2C - M_1\omega^2 & -C[1 + \exp(-iKa)] \\ -C[1 + \exp(iKa)] & 2C - M_2\omega^2 \end{vmatrix} = 0, \quad (21)$$

$$M_1M_2\omega^4 - 2C(M_1 + M_2)\omega^2 + 2C^2(1 - \cos Ka) = 0. \quad (22)$$

At $Ka \ll 1$, at the zone center

$$\omega^2 \cong 2C \left(\frac{1}{M_1} + \frac{1}{M_2} \right) \quad (\text{optical branch}); \quad (23)$$

Nearly a constant with K

$$\omega^2 \cong \frac{\frac{1}{2}C}{M_1 + M_2} K^2 a^2 \quad (\text{acoustical branch}). \quad (24)$$

Nearly linear with K

At $Ka = \rho, -\rho$ at the zone boundary

$$\omega^2 = 2C/M_1; \quad \omega^2 = 2C/M_2. \quad (25)$$

Optical and Acoustic Branches of the Dispersion for a Diatomic Linear Lattice




Figure 7 Optical and acoustical branches of the dispersion relation for a diatomic linear lattice, showing the limiting frequencies at $K = 0$ and $K = K_{\max} = \pi/a$. The lattice constant is a .

[111] Phonon Dispersion in Ge




Figure 8a Phonon dispersion relations in the [111] direction in germanium at 80 K. The two **TA + LA** phonon branches are horizontal at the zone boundary position, $K_{\text{max}} = (2\pi/a)(\frac{1}{2}\frac{1}{2}\frac{1}{2})$. The LO and TO branches coincide at $K = 0$; this also is a consequence of the crystal symmetry of Ge. The results were obtained with neutron inelastic scattering by G. Nilsson and G. Nelin.

[111] Phonon Dispersion in KBr




Figure 8b Dispersion curves in the [111] direction in KBr at 90 K, after A. D. B. Woods, B. N. Brockhouse, R. A. Cowley, and W. Cochran. The extrapolation to $K = 0$ of the TO, LO branches are called ω_T , ω_L .

Transverse Optical and Transverse Acoustic Waves of a Diatomic Linear Lattice


Substituting Eq. 23 to Eq. 20, we get

For $K = 0$, optical branch

$$M_1 u + M_2 v = 0$$

$$\frac{u}{v} = -\frac{M_2}{M_1} \quad (26)$$

Center of mass is fixed like a dipole as easily excited by E field in the optical wave.



For $K = 0$, acoustic branch, $u = v$

The atoms move in phase like acoustic wave in long wavelength.




Figure 10 Transverse optical and transverse acoustical waves in a diatomic linear lattice, illustrated by the particle displacements for the two modes at the same wavelength.

Quantization of Elastic Waves

The quantum of lattice vibration energy is called **phonon**, and the quantum number is denoted as **n** . The elastic waves in crystals are made of phonons.

$$\epsilon = (n + \frac{1}{2})\hbar\omega \quad (27)$$


$u = u_0 \cos Kx \cos \omega t$ for a standing wave

The time average kinetic energy is

$$\frac{1}{8} \rho V \omega^2 u_0^2 = \frac{1}{2} (n + \frac{1}{2}) \hbar \omega \quad , \quad (28)$$

$$u_0^2 = 4(n + \frac{1}{2}) \hbar / \rho V \omega \quad . \quad (29)$$

The sign of ω is usually positive; for imaginary ω , the crystal is unstable. An optical mode with ω close to zero is called a soft mode.



Phonon Momentum

Physical momentum of a crystal is

$$\mathbf{p} = M (d/dt) \sum_s u_s \quad (30)$$

$$\begin{aligned} \mathbf{p} &= M (du/dt) \sum_s \exp(is\mathbf{K}a) = \\ &M (du/dt) [1 - \exp(iN\mathbf{K}a)] / [1 - \exp(i\mathbf{K}a)] \end{aligned} \quad (31)$$


$$\sum_{s=0}^{N-1} x^s = (1 - x^N) / (1 - x) \quad (32)$$

For $\mathbf{K} = \pm 2\pi r/Na$, $\exp(iN\mathbf{K}a) = \exp(\pm i 2\pi r) = 1$

$$\mathbf{p} = M (du/dt) \sum_s \exp(is\mathbf{K}a) = 0 \quad (33)$$

The physical momentum of a crystal is zero.

Phonon Momentum



Elastic scattering of photons by a crystal

$$\mathbf{k}' = \mathbf{k} + \mathbf{G} , \quad (34)$$

For inelastic photon scattering, it creates a phonon momentum $\mathbf{K}$

$$\mathbf{k}' + \mathbf{K} = \mathbf{k} + \mathbf{G} . \quad (35)$$

For absorption of a phonon $\mathbf{K}$

$$\mathbf{k}' = \mathbf{k} + \mathbf{K} + \mathbf{G} . \quad (36)$$

Inelastic **neutron** scattering by phonons to obtain $\omega(\mathbf{K})$

$$\mathbf{k} + \mathbf{G} = \mathbf{k}' \pm \mathbf{K} , \quad (37)$$

$$\frac{\hbar^2 k^2}{2M_n} = \frac{\hbar^2 k'^2}{2M_n} \pm \hbar\omega , \quad (38)$$

Phonon Dispersions of Na in 3-D




Figure 11 The dispersion curves of sodium for phonons propagating in the [001], [110], and [111] directions at 90 K, as determined by inelastic scattering of neutrons, by Woods, Brockhouse, March and Bowers.


SUMMARY

- The quantum unit of a crystal vibration is a phonon. If the angular frequency is ω , the energy of the phonon is $\hbar\omega$.
- When a phonon of wavevector $\mathbf{K}$ is created by the inelastic scattering of a photon or neutron from wavevector $\mathbf{k}$ to $\mathbf{k}'$, the wavevector selection rule that governs the process is

$$\mathbf{k} = \mathbf{k}' + \mathbf{K} + \mathbf{G} ,$$

where $\mathbf{G}$ is a reciprocal lattice vector.

- All elastic waves can be described by wavevectors that lie within the first Brillouin zone in reciprocal space.
- If there are p atoms in the primitive cell, the phonon dispersion relation will have 3 acoustical phonon branches and $3p - 3$ optical phonon branches.



Chapter 4

- Problem set
 - No. 1, 3, and 4.
- Due 11/2, Wed. class